

(ii) *Dogs and cats*: It is used in certain acute and chronic dermatoses of varying etiology to help control the pruritus, irritation, and inflammation associated with these conditions.

(3) *Limitations*. Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infection. Do not use in animals with tuberculosis, chronic nephritis, cushingoid syndrome, or where peptic ulcers occur, except for emergency therapy. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 7131, Feb. 6, 1979, as amended at 61 FR 5506, Feb. 13, 1996]

§ 520.970 Flunixin oral dosage forms.

§ 520.970a Flunixin meglumine granules.

(a) *Specifications*. Each 10-gram packet contains flunixin meglumine equivalent to 250 milligrams of flunixin.

(b) *Sponsor*. No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 0.5 milligram of flunixin per pound of body weight (one packet per 500 pounds) per day.

(2) *Indications for use*. For alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

(3) *Limitations*. Administer daily dose for up to 5 days by sprinkling on small amount of feed. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 36381, June 22, 1979. Redesignated at 50 FR 38114, Sept. 20, 1985, and amended at 52 FR 7832, Mar. 13, 1987]

§ 520.970b Flunixin meglumine paste.

(a) *Specifications*. Each 30-gram syringe contains flunixin meglumine

equivalent to 1,500 milligrams of flunixin.

(b) *Sponsor*. No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Horses*—(1) *Amount*. 0.5 milligram of flunixin per pound of body weight daily.

(2) *Indications for use*. For alleviation of inflammation and pain associated with musculoskeletal disorders.

(3) *Limitations*. For oral use only. Treatment should not exceed 5 consecutive days. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 38114, Sept. 20, 1985, as amended at 52 FR 7832, Mar. 13, 1987]

§ 520.1010 Furosemide oral dosage forms.

§ 520.1010a Furosemide tablets or boluses.

(a) *Specifications*. Each tablet contains 12.5 or 50 milligrams of furosemide. Each bolus contains 2 grams of furosemide.

(b) *Sponsor*. See No. 012799 in § 510.600(c) of this chapter for conditions of use provided for in paragraphs (c) (1) and (2) of this section; see No. 000010 in § 510.600(c) of this chapter for use in dogs as provided for in paragraph (c)(1) of this section; see No. 000093 in § 510.600(c) of this chapter for use of a 12.5- and 50-milligram tablet for conditions of use provided for in paragraph (c)(3) of this section.

(c) *Conditions of use*. It is used as follows:

(1) *Dogs and cats*—(i) *Amount*. 1 to 2 milligrams per pound of body weight, once or twice daily, with a 6- to 8-hour interval between successive daily doses.

(ii) *Indications for use*. It is used for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency and acute non-inflammatory tissue edema.

(iii) *Limitations*. The dosage should be adjusted to the animal's response. In severe edematous or refractory cases, the dosage may be doubled or increased by increments of 1 milligram per pound